

EOS MLS Observations of the Two-Day Wave during 2004-2005

Varavut Limpasuvan

Department of Chemistry and Physics, Coastal Carolina University, Conway, South Carolina

Dong L. Wu, Michael J. Schwartz, and Joe W. Waters

NASA Jet Propulsion Laboratory, California Institute of Technology, Pasadena, California

Qian Wu and Tim L. Killeen

High Altitude Observatory, National Center for Atmospheric Research, Boulder, Colorado

Abstract. Two-day wave observations during January-March 2005 are reported using the recently launched Microwave Limb Sounder (MLS) aboard NASA's Earth Observing System Aura mission. Wave-induced disturbances in temperature, water vapor, carbon monoxide, and MLS line-of-sight wind appear in early January, peak near the end of January, and persist until late February. Temperature and wind amplitudes as large as 9 K and 50 m/s are observed near 90 km. The wave disturbance is initially confined in the mid to low summer latitudes where the climatological summer easterly jet exhibits strong shear. The wave then develops features akin to the third Rossby-gravity global normal mode, with a weak temperature disturbance in the winter hemisphere (anti-symmetric about the equator) and wind disturbance over the equator.

1. Introduction

Readily observed after the solstice above 40 km, the two-day wave is one of the most remarkable recurring features in the terrestrial atmosphere. The phenomenon is dominated by a westward propagating disturbance of zonal wavenumber 3 with a period of about 2.1 days, (3, 2.1) [Andrews et al., 1987]. It is also accompanied by smaller variances with wavenumber and period combinations, like (1, 6.7), (2, 3.5) and (4, 1.8), that tend to have similar phase speed (60-70 m/s) [Burks and Leovy, 1986; Garcia et al., 2005]. As such, the two-day wave appears to be an instability outgrowth of the mesospheric summer easterly jet [Plumb, 1984]. However, the wave also exhibits characteristics of an amplifying global normal mode under suitable supporting wind conditions [Salby, 1981; Randel, 1994].

Recent satellite observations report that wave-induced perturbations tend to be largest above the stratopause (~50 km). Near the upper mesosphere, temperature wave amplitude can exceed 5 K in the mid-latitude summer hemisphere [e.g. Garcia et al., 2005]. Two-day water vapor variation larger than 0.3 parts per million by volume (ppmv) can exist in regions of strong meridional gradient in water vapor distribution [Limpasuvan and Wu, 2003]. In low latitudes, meridional and zonal wind perturbations contribute significantly to the mesospheric flow, with maximum observed wave amplitude of nearly 60 and 20 m/s, respectively [Wu et al., 1993].

Here, the most recent two-day wave episode is reported using observations by Earth Observing System Microwave Limb Sounder (EOS MLS) aboard NASA's Aura satellite, launched in July 2004 [Waters et al., 2005]. In addition to two-day wave variations in temperature and water vapor, the present study provides a first glimpse at

the wave signature in carbon monoxide and in an offline-retrieved MLS wind product. As such, this study provides further evidence and insight of this remarkable phenomenon, whose global structures in the mesosphere/lower thermosphere are becoming more into focus with new measurements from NASA's Aura and TIMED (Thermosphere-Ionosphere-Mesosphere Energetics and Dynamics) spacecrafts. Additionally, as a recurring phenomenon, the two-day wave presence serves to further validate the wind observing techniques in the mesosphere.

2. Data and Analysis

This study uses EOS MLS temperature, water vapor (H_2O), and carbon monoxide (CO) profiles from an offline research retrieval for the period during December 2004 through March 2005. The data product spans nearly the entire globe with useful vertical range of 300-0.001 hPa for temperature and CO , and 300-0.1 hPa for H_2O . Vertical resolution for temperature is ~ 4 km in the middle stratosphere, decreasing to ~ 10 km above 0.1 hPa. For CO and H_2O , the vertical resolution is ~ 3 km.

Additionally, this study uses preliminary retrievals of mesospheric line-of-sight (LOS) wind from EOS MLS in the spacecraft's moving direction. The wind retrieval is based on the Doppler shift of the 118-GHz Zeeman-split O_2 lines, which can be resolved with 128 100-kHz-wide channels [Schwartz et al., 2005; Wu et al., 2005]. Reported here for the first time, the MLS LOS wind is a special MLS product, now under development. It typically has a (1/6-second integration time) single-measurement precision of ~ 15 m/s at 83-92 km to ~ 70 m/s at 70 km, which is the height range used in this study. For the polar-orbiting Aura satellite, the MLS LOS wind is dominated by the meridional component in the tropics, and therefore, winds derived from the northward (ascending)

and southward (descending) tracks are out-of-phase. While only the ascending LOS wind is presented in this paper, results from descending LOS wind are very similar.

To extract the two-day wave, the least-squared-fitting method is applied to the data [Wu et al., 1993; 1996], which has been successfully used to identify the two-day wave in the Upper Atmosphere Research Satellite (UARS) measurements. Based on an *a priori* assumption of the two-day wave recurring presence, the along-track profiles at each pressure levels are regressed onto a linear combination of cosine and sine functions of zonal wavenumber 3 of 0.48 cycles per day (~ 2.1 day period).

3. Results

Figure 1 (left) illustrates the evolution of the temperature two-day wave amplitude at 3 different pressure levels. A well-defined wave disturbance appears in the summer (Southern) hemisphere in early January and persists until near the end of February. At each level, peak amplitude exits around January 28, with largest amplitude (~ 9 K) found at the highest level.

The evolution of the two-day wave signature in the LOS wind differs from that in temperature (Figure 1, right). At each level, the primary peak amplitude occurs about 9 days earlier (January 19) in the mid summer latitudes. Then, around January 28, as the temperature wave matures, a weaker secondary peak in the LOS wind occurs with significant extension into the equatorial region at higher levels.

In the period of strongest temperature wave activity (Jan. 24-27), the cross-section of the temperature wave exhibits an inverted wedge-like shape in amplitude that ranges from 50 km to the upper observational limit (Figure 2a). With increasing altitude, the axis of maximum amplitude leans toward the summer pole as the amplitude latitudinal extent

grows (see also Figure 1). The wave phase tilts westward (overlaid black contours in Figure 2a) and the wave appears to have a very long vertical wavelength (> 40 km). A relatively weaker wave signature is evident in the winter (Northern) hemisphere near the stratopause above 80 km. The associated disturbances are out-of-phase with those found in the summer hemisphere, and are also present in Figure 1 (top-left).

Overall, the described characteristics and magnitude are strikingly similar to temperature observations discussed by Limpasuvan and Wu [2003] using UARS MLS data and by Garcia et al. [2005] using the TIMED Sounding of the Atmosphere using Broadband Emission Radiometry (SABER) data.

Cross-section of the LOS wind amplitude during January 24-27 shows a strong peak value of ~ 40 m/s near 40°S at 90 km (Figure 2b). A relatively weaker peak appears over the equatorial region at similar altitude (see also Figure 1, right). Like the temperature structure, the overall double-peak amplitude takes on an inverted wedge shape and the wave tilts westward with height (increasing phase line with altitude).

As the two-day wave signature in the meridional wind dominates over the zonal wind, the LOS wind wave amplitude should be dominated by the meridional wind disturbance [see also Section 2]. Indeed, the LOS wind amplitude structure and values are comparable to satellite observations by Wu et al. [1993], local station measurements [e.g. Fritts et al., 1999], and numerical model calculations [e.g. Palo et al., 1999]. Likewise, the shape is analogous to the third Rossby-gravity global normal mode meridional wind structure during solstice condition [Salby, 1981].

Strong wind perturbations induced by the two-day wave can elicit a strong response in density, thereby affecting tracer species in regions of strong meridional gra-

dient in tracer distribution [Limpasuvan and Wu, 2003]. Figure 3 demonstrates the concurrent evolution of the two-day wave amplitudes in H₂O and CO. Largely confined in the summer hemisphere, the H₂O maximum amplitude of ~0.42 ppmv exist near 45°S and 56 km and is nearly simultaneous with the temperature wave maximum [see Figure 1]. The evolution of CO tends to parallel the LOS wind response, peaking around 50°S prior to January 22 then at lower summer latitude around January 25. The maximum CO amplitude (~2.4 ppmv) is found above 90 km.

4. Summary and Discussion

This study presents the first observations of the two-day wave in temperature, water vapor, carbon monoxide, and LOS wind during December 2004-March 2005. The wave appears in early January, peaks near the end of January, and persists until late February. The temperature and wind amplitudes as large as 9 K and 50 m/s are observed near 90 km.

Pronounced wave amplitude appears at mid to low summer latitude in the region where the summer easterly jet tends to have strong vertical and meridional shear [e.g. CIRA; Fleming et al., 1990]. As such, wave growth from jet instability is a possible generating mechanism. However, the anti-symmetric (about the equator) temperature amplitude in the winter hemisphere, the simultaneous presence of equatorial wind disturbance, and the apparent large vertical wavelength are consistent with the third Rossby-gravity global normal mode [Salby, 1981; Garcia et al., 2005]. Thus, the observed two-day wave is consistent with previous interpretation that the wave arises from the excitation of normal modes by instability outgrowth of the easterly jet [Randel, 1994; Norton and Thuburn, 1996; Salby and Callaghan, 2001; Garcia et al., 2005].

Near the upper observational level, the evolution of wind and temperature amplitude suggests the normal mode excitation process (Figure 1, top). The early peaking of the LOS wind amplitude over a deep layer in the mid summer latitude (also mimicked by CO; Figure 3) may be related to instability growth prior to January 22. This growth is followed by the characteristic wave features of the third Rossby-gravity global normal mode around Jan. 25-27, with winter temperature disturbance and equatorial wind enhancement.

In addition, during January 16-22, a strong solar proton event was observed with numerous medium and large solar flares from a single sunspot cluster. This event may have considerable impact on the mesosphere dynamics [e.g. Seppala et al., 2005]. On January 22, the NOAA GOES-11 observed an in-flux of solar protons (> 100 MeV) to be 6.1×10^6 Protons/cm²-day-sr, on the same order as the famous October 1989 solar proton event. Roles of upper-air instability and strong solar activity in the 2-day wave excitation warrant further investigations.

The LOS wind product is preliminary, but comparisons with concurrent TIMED Doppler Interferometer (TIDI; Killeen et al., 1999; Skinner et al., 2003) measurements appear very favorable. By projecting TIDI horizontal wind components (using the cold-side profiles along the ascending tracks) onto the MLS LOS, resulting LOS wind wave exhibits structure, amplitude, and phase similar to those observed by MLS (compare Figure 2b and 2c). In fact, the LOS wind amplitude in Figure 2c greatly resembles the two-day wave in TIDI *meridional* wind below 105 km (not shown). Finally, at 90 km, reconstructed wave oscillation in LOS winds demonstrates a strong one-to-one correlation throughout the entire globe between TIDI and MLS (Figure 4).

Acknowledgements. V. L. is supported by the NASA and South Carolina Space Grant and CCU's Public Engagement Program. Part of this work was performed at the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration.

References

- Andrews, D. G., J. R. Holton, and C. B. Leovy, 1987: *Middle Atmospheric Dynamics*, Academic Press, 489 pp.
- Burks, D. and C. B. Leovy, 1986: Planetary waves near the mesospheric easterly jet, *Geophys. Res. Lett.*, **13**, 193-196.
- Fleming, E. L. and co-authors, 1990: Zonal mean temperature, pressure, zonal wind, and geopotential height as function of latitude, *Adv. Space Res.*, 10, Issue **12**, 11-59.
- Fritts, D. C. and co-authors, 1999: Two-day wave structure and mean flow interactions observed by radar and High Resolution Doppler Imager, *J. Geophys. Res.*, **104**, 3953-3969.
- Garcia, R. R. and co-authors, 2005: Large-scale waves in the mesosphere and lower thermosphere observed by SABER, *J. Atmos. Sci.*, in review.
- Skinner, W. R. and co-authors, 2003: Operational Performance of the TIMED Doppler Interferometer (TIDI), *Proc. SPIE*, 5157, 47-57.
- Killeen, T. L. and co-author, 1999: TIMED Doppler Interferometer (TIDI), *Proc. SPIE* 3756, 289-301, *Optical Spectroscopic Techniques and Instrumentation for Atmospheric and Space Research III*, Allen M. Larar Ed.
- Limpasuvan, V. and D. L. Wu, 2003: Two-day wave observations of UARS Microwave Limb Sounder mesospheric water vapor and temperature, *J. of Geophys. Res.*, **108**, No. D10, 4307, doi: 10.1029/2002JD002903.
- Norton, W. A. and J. Thuburn, 1996: The two-day wave in the middle atmosphere GCM, *Geophys. Res. Lett.*, **23**, 2113-2116.
- Palo, S. E., R. G. Roble, and M. E. Hagan, 1999: Middle atmosphere effects of the quasi-two-day wave determined from a General Circulation Model, *Earth Planets Space*, **51**, 629-647.
- Plumb, R. A., 1983: Baroclinic instability at the Summer Mesosphere: a mechanism for the quasi-two-day wave?, *J. Atmos. Sci.*, **40**, 262-270.

- Randel, W. J., 1994: Observations of the 2-day wave in NMC stratospheric analyses, *J. Atmos. Sci.*, **51**, 306-313.
- Salby, M. L., 1981: Rossby normal modes in nonuniform background conditions. Part II: Equinox and solstice conditions, *J. Atmos. Sci.*, **38**, 1827-1840.
- Salby, M. L. and P.F. Callaghan, 2001: Seasonal Amplification of the 2-Day Wave: Relationship between Normal Mode and Instability, *J. Atmos. Sci.*, **58**, 1858-1869.
- Schwartz, M. J. and co-authors, 2005: MLS forward model for polarized mesospheric signals, *IEEE Trans. Geos. & Remote Sens.*, in review.
- Seppala, A. and co-authors, 2005: The high energy proton fluxes of January 2005: Impact on middle atmosphere, *to be presented* at International Association of Geomagnetism and Aeronomy (IAGA) 2005 Scientific Assembly (Toulouse, France).
- Waters, J. W. and co-authors, 2005: The Earth Observing System Microwave Limb Sounder (EOS MLS) on the Aura Satellite, *IEEE Trans. Geos. & Remote Sens.*, in review.
- Wu, D. L. and co-authors, 1993: Observations of the quasi 2-day wave from the High Resolution Doppler Image on UARS, *Geophys. Res. Lett.*, **24**, 2853-2856.
- Wu, D. L. and co-authors, 1996: Excitation and evolution of the quasi-2-day wave observed in UARS/MLS temperature measurements, *J. Atmos. Sci.*, **53**, 728-738.
- Wu, D. L. and M. J. Schwartz, 2005: Mesospheric wind measurements from EOS MLS 118GHz radiometer, *in preparation*.

V. Limpasuvan, Coastal Carolina University, P. O. Box 261954, Conway, SC 29528, USA. (E-mail: var@coastal.edu)

D. L. Wu, J. W. Waters, and M. J. Schwartz, NASA JPL, Caltech, 4800 Oak Grove Drive, Pasadena, CA 91109, USA. (E-mail: dwu, joe, or michael@mls.jpl.nasa.gov)

Q. Wu and T. L. Killeen, High Altitude Observatory, NCAR, P.O. Box 3000, Boulder, CO 80307, USA (E-mail: qwu or killeen@ucar.edu)

LIMPASUVAN ET AL.: EOS MLS TWO-DAY WAVE 2004-2005


FIGURES AND CAPTIONS

Figure 1. The 2005 latitude-time evolution of the two-day wave amplitude in temperature (every 2 K) and line-of-sight (LOS) wind (every 5 m/s). The LOS wind is computed from profiles along the ascending tracks (see Section 2).




Figure 2. January 24-27 meridional structure of the two-day wave amplitude (filled contours) of T (a, in K) and LOS wind (b, in m/s) from ascending MLS data, and ascending cold side TIDI profiles (c, in m/s). Phase lines (every 45°) are in black.




Figure 3. The 2005 latitude-time evolution of the two-day wave amplitude in carbon monoxide (top; every 0.5 ppmv) and water vapor (bottom; every 0.2 ppmv)




Figure 4. Simultaneous correlation between LOS wind two-day wave perturbation maps from TIDI and MLS. Each star represents values of common region on the globe at 90 km.